

Section 17.1

Ottoson D, Shepherd GM (1970). Steps in impulse generation in isolated muscle spindle. *Acta Physiologica Scandinavica* 79: 423-430.

Section 17.2

Bakken GS, Krochmal AR (2007). The imaging properties and sensitivity of the facial pits of pitvipers as determined by optical and heat-transfer analysis. *Journal of Experimental Biology* 210: 2801–2810.

Baylor DA, Fuortes MG, O'Brien PM (1971). Receptive fields of cones in the retina of the turtle. *Journal of Physiology* 214: 265-294.

Baylor DA, Lamb TD, Yau KW (1979). The membrane current of single rod outer segments. *Journal of Physiology* 288: 589–611.

Fesenko EE, Kolesnikov SS, Lyubarsky AL (1985). Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. *Nature* 313: 310–313.

Fuortes MGF 1959. Initiation of impulses in visual cells of *Limulus*. *Journal of Physiology* 148: 14-28.

Konopka RJ, Benzer S (1971). Clock mutants of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America* 68: 2112–2116.

Mäthger LM, Roberts SB, Hanlon RT (2010). Evidence for distributed light sensing in the skin of cuttlefish, *Sepia officinalis*. *Biology Letters* 6: 600–603.

Merlin C, Gegear RJ, Reppert SM (2009). Antennal circadian clocks coordinate sun compass orientation in migratory monarch butterflies. *Science* 325: 1700-1704.

Patek SN, Caldwell RL (2005). Extreme impact and cavitation forces of a biological hammer: strike forces of the peacock mantis shrimp. *Journal of Experimental Biology* 208: 3655-3664.

Vitaterna MH, King DP, Chang AM, Kornhauser JM, Lowrey PL, McDonald JD, Dove WF, Pinto LH, Turek FW, Takahashi JS (1994). Mutagenesis and mapping of a mouse gene, *Clock*, essential for circadian behavior. *Science* 264: 719–725.

Zhu H, Yuan Q, Briscoe AD, Froy O, Casselman A and Reppert SM (2005). The two CRYs of the butterfly. *Current Biology* 15: R953-R954.

Section 17.3

Bushdid C, Magnasco MO, Vosshall LB, Keller A (2014). Humans can discriminate more than 1 trillion olfactory stimuli. *Science* 343: 1370-1372.

Caprio J, Brand JG, Teeter JH, Valentincic T, Kalinoski DL, Kohbara J, Kumazawa T, Wegert S (1993). The taste system of the channel catfish: from biophysics to behaviour. *Trends in Neurosciences* 16: 192-197.

Fine JM, Sorensen PW (2005). Biologically-relevant concentrations of petromyzonol sulphate, a component of the sea lamprey migratory pheromone, measured in stream waters. *Journal of Chemical Ecology* 31: 2205-2210.

Freeman EG, Wisotsky Z, Dahanukar A (2014). Detection of sweet tastants by a conserved group of insect gustatory receptors. *Proceedings of the National Academy of Sciences of the United States of America* 111: 1598-1603.

Gamal AM, Elsheikh EH, Nasr ES (2012). Morphological adaptation of the buccal cavity in relation to feeding habits of the omnivorous fish *Clarias gariepinus* : A scanning electron microscopic study. *The Journal of Basic and Applied Zoology* 65: 191 – 198.

Green PA, Valkenburgh B, Pang B, Bird D, Rowe T, Curtis A (2012). Respiratory and olfactory turbinal size in canid and arctoid carnivorans. *Journal of Anatomy* 221: 609-621.

Royer SM, Kinnamon JC (1991). HVEM serial-section analysis of rabbit foliate taste buds. I. Type III cells and their synapses. *Journal of Comparative Neurology* 306: 49-72.

Section 17.4

Christensen CB, Christensen-Dalsgaard J, Brandt C, Madsen PT (2012). Hearing with an atympanic ear: good vibration and poor sound-pressure detection in the royal python, *Python regius*. *Journal of Experimental Biology* 215: 331-342.

Langbauer WR, Payne KB, Charif RA, Rapaport L, Osborn F (1991). African elephants respond to distant playbacks of low-frequency conspecific calls. *Journal of Experimental Biology* 157: 35-46.

Li W, Feng Z, Sternberg PW, Shawn Xu XZ (2006). A *C. elegans* stretch receptor neuron revealed by a mechanosensitive TRP channel homologue. *Nature* 440: 684–687.

Pawson L, Prestia LT, Mahoney GK, Güçlü B, Cox PJ, Pack AK (2009). GABAergic/glutamatergic-glial/neuronal interaction contributes to rapid adaptation in Pacinian corpuscles. *Journal of Neuroscience* 29: 2695–2705.

Ricci AJ, Bai J-P, Song L, Lv C, Zenisek D, Santos-Sacchi J (2013). Patch-clamp recordings from lateral line neuromast hair cells of the living zebrafish. *Journal of Neuroscience* 33: 3131–3134.

Shin J-B, Adams D, Paukert M, Siba M, Sidi S, Levin M, Gillespie PG, Gründer S (2005). Xenopus TRPN1 (NOMPC) localizes to microtubule-based cilia in epithelial cells, including inner-ear hair cells. *Proceedings of the National Academy of Sciences of the United States of America* 102: 12572–12577.

Section 17.5

Gracheva EO, Ingolia NT, Kelly YM, Cordero-Morales JF, Hollopeter G, Chesler AT, Sánchez EE, Perez JC, Weissman JS, Julius D (2010). Molecular basis of infrared detection by snakes. *Nature* 464: 1006-1011.

Saito S, Shingai R (2006). Evolution of thermoTRP ion channel homologs in vertebrates. *Physiological Genomics* 27: 219-230.

Sidi S, Friedrich RW, Nicolson T (2003). NompC TRP channel required for vertebrate sensory hair cell mechanotransduction. *Science* 301: 96–99.

Section 17.6

Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, Koltzenburg M, Basbaum AI, Julius D (2000). Impaired nociception and pain sensation in mice lacking capsaicin receptor. *Science* 288: 306-313.

Tracey WD Jr, Wilson RI, Laurent G, Benzer S (2003). Painless, a *Drosophila* gene essential for nociception. *Cell* 113: 261-273.

Summers T, Holec S, Burrell BD (2014). Physiological and behavioral evidence of a capsaicin-sensitive TRPV-like channel in the medicinal leech. *Journal of Experimental Biology* 217: 4167-4173.

Section 17.7

Clarke D, Whitney H, Sutton G, Robert D (2013). Detection and learning of floral electric fields by bumblebees. *Science* 340: 66–69.

Czech-Damal NU, Liebschner A, Miersch L, Klauer G, Hanke FD, Marshall C, Dehnhardt G, Hanke W (2012). Electroreception in the Guiana dolphin (*Sotalia guianensis*). Proceedings of Royal Society B 279: 663–668.

Jackson CW, Hunt E, Sharkh S, Newland PL (2011). Static electric fields modify the locomotory behaviour of cockroaches. Journal of Experimental Biology 214: 2020–2026.

Proske U, Gregory JE, Iggo A (1998). Sensory receptors in monotremes. Philosophical Transactions of the Royal Society B 353: 1187–1198.

von der Emde G, Fetz, S (2007). Distance, shape and more: recognition of object features during active electrolocation in a weakly electric fish. Journal of Experimental Biology 210: 3082–3095.

Section 17.8

Engels S, Schneider N-L, Lefeldt N, Hein CM, Zapka M, Michalik A, Elbers D, Kittel A, Hore PJ, Mouritsen H (2014). Anthropogenic electromagnetic noise disrupts magnetic compass orientation in a migratory bird. Nature 509: 353–356.

Gegear RJ, Casselman A, Waddell S, Reppert SM (2008). Cryptochrome mediates light-dependent magnetosensitivity in *Drosophila*. Nature 454: 1014–1018.

Freire R, Eastwood MA, Joyce M (2011). Minor beak trimming in chickens leads to loss of mechanoreception and magnetoreception. Journal of Animal Science 89: 1201–1206.